

24th Annual

Cardiologists Conference

June 11-13, 2018 | Barcelona, Spain



Damien Byas

Center for Healthcare and Organizational Research, USA

Examining risk factors associated with worldwide sudden cardiac death rates in children and adults

Statement of the Problem: Physical inactivity, diabetes, diets high in fats and cholesterol, and obesity are all considered as poor health conditions which serve as significant gateway factors which may lead to heart disease and possibly cardiac death. In 2012 worldwide (World Health Federation, 2016), there were more than 17.3 million cardiac related deaths. Cardiovascular diseases were the main cause of death in almost all Organization for Economic Co-operation and Development (OECD) countries, and accounted for 35% of all deaths in 2009 (OECD, 2011). This study examined specific identifiable risk factors which may be associated with sudden cardiac death rates in children and adult populations using the Kids' Inpatient Database (KID), Healthcare Cost and Utilization Project (HCUP), and the Agency for Healthcare Research and Quality (AHRQ, 2016).

Orientation: A large randomly drawn sample (N=422,599) of boys (n=198,960) and girls (n=223,639) ages 4 to 12, was examined in this research study to test for the association between risk factors which may be associated with heart disease prevalence. The Pearson Chi Square test was applied to measure for significant variable relationships in this research study.

Methodology & Theoretical Orientation: The Pearson chi square test was applied to measure for significant variable relationships in this research study.

Findings: The results of this study found that there was a statistically significant association between cardiovascular disease prevalence and identifiable risk factors in children and adult groups ($p < .05$). Other significant associations were also found as a result of the Chi square analysis.

Conclusion & Significance: Recommendations are made to implement more effective strategies for health promotion and disease prevention, health education, cultural competence training for healthcare professionals, improve healthcare quality, and economic development.

Levels of Association	Strength of Association
< .10	Weak
.11 to .30	Moderate
.31 to .35	Strong
> .35	Very Strong

Standard for Cramer's V and Phi coefficients

24th Annual

Cardiologists Conference

June 11-13, 2018 | Barcelona, Spain

Recent Publications

1. Byas D K, Gillespie J, Talbert P Y and Kinnard J (2016) Examining the association between healthcare quality and health status among American children. *Journal of the National Society of Allied Health*, Fall/Winter 2016 edition. 13(1): 25-29.
2. Byas D (2011) Developing strategies to reduce children's health care disparities. *Internet Journal of Healthcare Administration* 7(2).

Biography

Damien Byas is currently serving as an Associate Faculty Member in a Master of Public Health (MPH) program; President of North American Scientific Committee on Cardiovascular Health; a part-time public health researcher, Senior Research Fellow, and an Adjunct Professor for an MPH program.

Notes: