

12th International Conference on

PHARMACEUTICAL CHEMISTRY

&

2nd EUROPEAN PATHOLOGY CONGRESS

May 20-21, 2019

Berlin, Germany

**Andrea M P Romani**

Case Western Reserve University, USA

Cellular magnesium as a regulator of glucose homeostasis and insulin-mediated signaling and cellular metabolism

The last thirty years have registered a progressive and dramatic increase in the incidence of obesity and type 2 diabetes mellitus world-wide. Metabolic Syndrome, one of the most commonly conditions associated with obesity and insulin resistance, has also increased considerably. The latest releases from the WHO estimate that approximately 1 billion people worldwide are obese, and more than 500 million are diabetic or pre-diabetic. Interestingly, increasing evidence suggest that our current western diet is hypercaloric but hyponutritive, as it is lacking essential micronutrients and minerals. Our laboratory has focused on the possible role of reduced cellular magnesium levels in the dysregulation of cellular and systemic glucose homeostasis. Experimental data obtained in animal and cellular models, including cells of human origin support the conclusion that cellular magnesium regulates transmembrane glucose transport as well as its utilization, and neosynthesis in gluconeogenic tissues, by modulating the activity of specific cellular enzymes and insulin-mediated signaling. Regardless of the tissue considered,

decreased cellular and serum magnesium levels impact the proper operation of Glut 4, and Glut 2 transporters, thus limiting the ability of tissues like heart, muscles, liver, and possibly beta-islets, to effectively transport glucose into the cell to support glycolysis, ATP production, and ultimately storage as glycogen. As a consequence, gluconeogenesis becomes erroneously activated, further enhancing the circulating levels of glucose and resulting in the dysregulation of fatty acids, cholesterol, and protein degradation, to support gluconeogenic activity through increased cortisol production and insulin resistance. Also, decreased cellular magnesium levels appear to contribute directly to increased basal inflammation within tissues, further impairing insulin responsiveness and systemic metabolic homeostasis. Altogether, our results argue for the necessity to better understand the role that micronutrients play in modulating both organ-specific and systemic metabolism and inflammation, to ultimately identifying more effective therapeutic and dietary approaches.

Biography

Andrea M P Romani, obtained his medical degree from the University of Siena, Italy and his PhD from the University of Turin, Italy. Upon completing his postdoctoral studies under Dr. Scarpa, he joined the faculty in the Department of Physiology and Biophysics, Case Western Reserve University, where he is currently the Associate Professor. He has published over 90 peer review articles in high profile journals together with numerous invited reviews and book chapters on the role of mammalian magnesium homeostasis in health and disease. He is currently serving as an Editorial Board Member and an ad hoc reviewer for numerous international journals including Biochemical Pharmacology and Plos ONE.

andrea.romani@case.edu